

Plant resistance to extreme temperatures

Plants have a reduced mobility and have no other choice than to face the variations of the physical conditions of their environment. In order to adjust and adapt to new conditions they must **perceive** the changes and **translate** them **into signals** capable of being directed to a **target** that will perform the operations that will allow them to maintain their activity. The response of plants to their environment therefore involves **receptors** and **signalling molecules**.

- The targets can be (1) the nuclear, chloroplast or mitochondrial **genome**, (2) **transcription factors**, or (3) transcription factor **activators**. Syntheses are then initiated or modulated, or even stopped. The response often affects both metabolism and morphogenesis.
- Note that a **signal** from a modification **perceived** only **at one point of the plant** (on one of its leaves, for example) **can be transmitted** to the whole plant, and provoke a global adaptive response that is called **systemic response**.

It is difficult to identify a single thermal receptor that would link the plant to its environment: the plant probably has several at key points of metabolism. However, it **seems necessary** that the signals they produce **interact** to produce the **appropriate global response**. Finally, it should be noted that the diversity of systems that "perceive" temperature changes can contribute to the diversity of the plant world.

Research using the possibilities of molecular genetics has developed in the last two decades. It has considerably deepened our knowledge of the response to extreme temperatures. But many points remain to be clarified. This focus presents a few avenues that will give the reader an idea of the complexity of the problems posed.

1. Hardening and light: the "phytochrome" thermometer

The interactions between light and cold hardening have long been highlighted (see [Effect of temperature on photosynthesis](#)). In some cases, cold hardening can even be obtained at ordinary temperature **by modulating the length and spectral composition of the photoperiod**. However, cold hardening is still necessary to obtain a complete hardening

Phytochromes (P) are a family of chromoproteins that allow the plant to perceive the photoperiod length and spectral composition in the red (Figure 1). They are well known to be involved in the processes of morphogenesis and flowering.

- Their inactive form (P_R) preferentially absorbs red light (R) to give an active form P_{FR} preferentially absorbing dark red (FR).
- **This reaction is reversible**: P_{FR} absorbing RS gives again the inactive form (P_R). The P_{FR}/P_R ratio in the plant exerts control over many plant responses to light (morphogenesis, flowering etc.) by controlling signalling pathways leading to modulation of gene activity.
- **But the switch from P_{FR} (active form) to P_R (inactive form) also occurs independently of light**. This reaction is strongly influenced by temperature. It can therefore modulate the P_{FR}/P_R ratio, and consequently the activity of phytochromes, independently of the characteristics of the light environment.
- It constitutes a **thermometer able to appreciate a difference of 1°C, at least**. [\[1\]](#)




Figure 1. Interconversion of the red light absorbing phytochrome P_R to the active dark red light absorbing phytochrome P_{FR} . P_{FR} also gives P_R in a reaction that is not light sensitive but is temperature dependent. [Source: Author's diagram]

P_{FR} , transferred into the cell nucleus, modulates cellular responses caused by low or high temperatures. This regulation occurs at the level of transcription factors that control, among other things, several aspects of morphogenesis [2].

2. Membrane fluidity: its relationship with related proteins

Biological membranes, lipid bilayers, are composed of a mixture of unsaturated fatty acids (with one or more double bonds), which are fluid, and saturated fatty acids (do not have double bonds), which are comparatively more rigid.

Their fluidity increases with temperature, affecting not only their permeability but also the activity of the proteins included in them or interacting with them.

This modification of activity can thus be the cause, by successive cascades, of a modulation of the activity of genes.

However, fluidity tends to be maintained, despite temperature variations, by a rapid modulation of the ratio between saturated and unsaturated fatty acids [3], with an influence on the activity of the proteins included in it.




Figure 2. Rapid temperature variations induce rapid changes in membrane fluidity. The latter is high when the temperature is high and low when it is low. The variation of the ratio (Saturated Lipids) / (Unsaturated Lipids) in plants growing under different thermal climates allows the adjustment of the membrane fluidity for an optimal functioning of the proteins included in it and the regulation of the compartment to compartment exchanges they control. [Source: Author's diagram]

It is also possible that the increase in membrane fluidity is at the origin of the influx of Ca^{++} into the cytoplasm. This influx affects the processes leading to **thermal tolerance**.

3. Isoprene synthesis

Isoprene is emitted into the atmosphere by many woody species (such as oaks, eucalyptus, conifers) and herbaceous species. Several species of mosses (such as *Sphagnum*) and ferns also produce isoprene. However, many plants do not emit isoprene: this is the case, for example, of the bean, although other papilionaceous plants produce it.

Isoprene **protects the photosynthetic apparatus** [4] against **thermal shock**. This protection is also effective in plants that emit

little or no isoprene, placed in an atmosphere that contains levels compatible with those measured near the leaves of emitting plants.

Isoprene is thought to contribute to the rapid **stabilization of membranes**, but also to the **regulation of the quantity** of active O₂ species. Finally, it acts as a **signalling molecule** by acting on all the systems involved in growth and in responses to stress.

4. The production of active oxygen species (ROS)

The production of active oxygen species is common in plants subjected to all kinds of constraints (physical or biological). It occurs, at both low and high temperatures, mainly in chloroplasts, mitochondria and peroxisomes.

For example, active oxygen species (ROS) [5] are **continuously produced during photosynthesis**. Indeed, as the absorption of light by the collecting antennae and its transfer to the reaction centers of the photosystems are insensitive to temperature, any decrease in the assimilation of CO₂ produced by extreme temperatures (low and high) will result in excess light energy in the photosystems (Read [How do plants cope with alpine stress?](#))

- **Part of this energy is evacuated in the form of heat** from the antennae;
- **Another part contributes to the formation of ROS**, which can **damage** the photosynthetic apparatus or **trigger acclimatisation** responses.

ROS are formed at other stages in the functioning of the system of transduction of light energy into chemical energy. They trigger adaptive responses by interacting with the metabolic and signalling networks already at work.

They also participate, probably with calcium, in **systemic responses**. Indeed, it has been shown that an **oxidative wave** arises at **the point of the plant** subjected to an increase in temperature and **spreads to the entire plant**. It originates from the production of ROS at the point affected by the stress (Figure 3). This propagation occurs from cell to cell in two ways:

1. By a **symplasmic pathway**, i.e. from cytoplasm to cytoplasm via plasmodesmata and channels, such as aquaporins;
2. By an **apoplastic pathway**, located between the skeletal wall and the plasmalemma. This last pathway involves an NADPH oxidase located in the plasmalemma whose activation requires Ca⁺⁺. A "**calcium wave**" accompanies the oxidative wave.




Figure 3. Diagram showing that a localized stress can trigger a global response of the plant allowing it to resist this stress. This is the systemic response. It originates from the production of ROS at the point affected by the stress. [Source: Author's diagram]

These signals are transmitted rapidly, probably through the vascular system. They lead to very rapid systemic responses at the molecular level: in *Arabidopsis thaliana*, for example, we have been able to show an ultra-rapid response of the genome (in a few seconds) with the appearance of transcripts of proteins associated with the response to abiotic stress [6].

Cover image. [Source: © G.Cornic]

- [1] Jung, J-H., Domijan, M., Klose, C., Biswas, S., Ezer, D., Gao, M., Khattak, A.K., Box, M.S., Charoensawan, V., Cortijo, S., Kumar, M., Grant, A., Locke, J.C.W., Schäfer, E., Jaeger, K.E., Wigge, P.A. (2016) Phytochromes function as thermosensors in *Arabidopsis*. *Science* 354, 886-889
- [2] Catalá, R., Medina, J., Salinas, J. (2011) Integration of low temperature and light signaling during cold acclimation response in *Arabidopsis*. *Proceedings of the National Academy of Sciences* 108, 39, 16475-16480
- [3] Sung DY, Kaplan F, Lee KJ, Guy CL (2003) Acquired tolerance to temperature extremes. *Trends in Plant Science* 8, 179-187
- [4] Sharkey TD, Chen X, Yeh S (2001) Isoprene increases thermotolerance of foscimidomycin-fed leaves. *Plant Physiology* **125**, 2001-2006
- [5] Reactive Oxygen Species = ROS. ROS are radicals that have a single electron on their peripheral layer, a characteristic that allows them to interact easily with their environment.
- [6] Suzuki N, Devireddy AR, Inupakutika MA, Baxter A, Mille G, Song L, Shulaev E, Rajeev K. Azad, Shulaev V, Mittler R (2015) Ultra-fast alterations in mRNA levels uncover multiple players in light stress acclimation in plants. *The Plant Journal* **84**, 760-772

L'Encyclopédie de l'environnement est publiée par l'Université Grenoble Alpes.

Les articles de l'Encyclopédie de l'environnement sont mis à disposition selon les termes de la licence Creative Commons Attribution - Pas d'Utilisation Commerciale - Pas de Modification 4.0 International.
